

# Hypoxic Challenge (Flight Assessment)

A patient's guide

### What is a Hypoxic Challenge?

When a person goes on a flight, there is less oxygen available in the air on board than normal. The *Hypoxic Challenge* (also known as a flight assessment test), helps us to understand how well a person's body will cope with this change in oxygen during a flight.

The results of this test will help doctors to decide whether someone with lung disease needs extra oxygen when flying.

#### Who needs the test?

If you are planning to fly, you may be asked to have a hypoxic challenge assessment if:

- you have a breathing disorder that causes you to carry less oxygen in your blood.
- you already use oxygen as part of your treatment.
- you have experienced a noticeable deterioration in your condition since you last went on a flight.

 you have previously had breathing problems or associated symptoms, during or following a flight.

#### What does the test involve?

Normally, there is 21% oxygen in the air we breathe, but during the test you will breathe air containing 15% oxygen. This is the same as the air you would breathe in an aircraft cabin.

The test is performed in the Respiratory Physiology department by a clinical physiologist. It consists of a maximum of three 20 minute test phases, with some baseline readings taken at the very start and recovery readings at the end.

- You will wear a probe on your forehead, which will monitor your oxygen levels and pulse rate throughout the test.
- You will wear an open, clear face mask with an attachment. This will be connected to a supply of gases, which lowers the level

of oxygen you are breathing to the required 15%.

 Underneath this mask, you will wear a nasal tube, which will deliver extra oxygen during the test, if it is required.

At the very beginning of the test and at the end of each 20 minute phase, you will have an earlobe *Capillary Blood Gas* test. This is to accurately measure your body's response to the changes in the level of oxygen.

- A small amount of Deep Heat cream will be put on your earlobe for 15 minutes. This will flush the area with blood.
- Once the cream is removed, the area is sterilised and a small scratch is made to draw a blood sample, which is then analysed immediately.
- At each stage, the result of this blood test determines if we need to increase the level of oxygen and continue.

### **Frequently Asked Questions**

### How long will the test take?

The length of the test depends upon how many phases we have to perform. The shortest test may take 45 minutes, where the longest test may take around 90 minutes.

We advise you to set aside up to 2 hours for the test, to allow for any delay.

## What do I need to bring with me?

Please bring sufficient change for car parking facilities (£3.50 for 4 hours / £5.00 per day).

It may be useful to bring a list of medications you are taking, or a copy of your prescription, so that we know what medication you are on at the time. There is no need to withhold any medication prior to this test.

As the test may take some time, with little activity on your part, you may want to bring a book or magazine to read, or music to listen to.

Please note - due to our location and the equipment we use, you may not be able to connect to the internet.

# What symptoms might I get during the test?

The hypoxic challenge rarely causes any side-effects.

On occasion some people may experience a dry mouth, chest tightness, drowsiness or slight headache. You will be able to tell your physiologist about any symptoms during the test. Depending on the nature of the symptom, the test may be stopped early. Such circumstances are rare and normally resolve rapidly upon breathing room air.

### How do I find out my results?

The clinical physiologist performing your test will not be able to give you your results. It is up to your consultant to make appropriate recommendations for your plans to travel. Depending on the urgency, this may take up to three weeks to achieve. In some circumstances you may

need to contact your consultant directly to discuss the result. Your physiologist will be able to provide you with the necessary contact details.

## Who do I contact if I need further information?

Please contact the Respiratory Physiology department if you have any questions or concerns about your test or appointment.

We can be reached by phone on 01480 364815 between 9am and 4.30pm.

### **Royal Papworth Hospital NHS Foundation Trust**

Papworth Everard, Cambridge, CB23 3RE

Tel: 01480 830541 Fax: 01480 831315

### www.royalpapworth.nhs.uk

A member of Cambridge University Health Partners



Royal Papworth Hospital is a smokefree site







Author ID: Clinical Physiologist
Department: Respiratory Physiology
Printed: October 2018

Review date: October 2020

Version:

Leaflet number: PI 196

Large print copies and alternative language versions of this leaflet can be made available on request.

© Royal Papworth Hospital NHS Foundation Trust